

Hackman, 1980

Notulae Entomologicae 60:117—162, 1980.

A check list of the Finnish Diptera II. Cyclorrhapha

Walter Hackman

Abstract

HACKMAN, WALTER: A check list of the Finnish Diptera II. Cyclorrhapha.
— Notulae Entomol. 60:117—162, 1980.

A check list of the Diptera found within the present borders of Finland.
Part II contains the Cyclorrhapha, which comprise 58 families and 2438
species. Corrections to part I and a generic index to parts I and II are
given.

Author's address: Prof. Walter Hackman, Zoological Museum, N. Järn-
vägsg. 13, SF-00100 Helsingfors 10.

The first part of this check list of
the Finnish Diptera was published
earlier this year in this journal (HACK-
MAN, 1980) and contained the Nema-
tocera and the Brachycera (s.str.). The
present list contains the Cyclorrhapha
and follows the same principles as the
first part; eastern Fennoscandian spe-
cies found only outside the present
borders of Finland have also been left
out of this part of the list.

The list covering the Pipunculidae
was compiled by Mr. Anders Albrecht
(Zoological Museum, Helsinki) and the
list covering the Platypezidae was
checked by Dr. Peter Chandler
(Weston Research laboratories, Mai-
denhead, England).

In general, the system presented by
STEYSKAL (1974) and based on the
work of GRIFFITHS (1972) has been
followed. There are, however, the
following exceptions: The Conopidae
have been placed after Syrphoidea, in
a superfamily of their own, Conopo-
idea and not included in Tephritoidea.
The Diastatidae are retained as a
family including the genera Campi-
choeta and Diastata. The Oestridae

and Gasterophilidae are placed in
Oestroidea. In Tachinoidea, the Rhi-
nophoridae, Sarcophagidae, Calliphoro-
idae and Phasidae are considered as
families separate from Tachinidae.
Hippoboscidae and Nycteribiidae are
kept as separate families in the super-
family Glossinoidea.

The genus *Ornitholeria* Frey is ac-
cording to Dr. J. F. Vockeroth (un-
published), a junior synonym of *Neos-
sos* Malloch and has been placed in
the Chiropteromyzidae as suggested to
me by Mr. Brian Cogan (British Mu-
seum of Nat. Hist., London).

References

- GRIFFITHS, G. C. D. 1972: The phylogenetic
classification of Diptera Cyclorrhapha,
with special reference to the structure
of the male post-abdomen. — 340 pp.
The Hague.
HACKMAN, W. 1980: A check list of the Fin-
nish Diptera I. Nematocera and Brachy-
cera (s. str.). — Notulae Entomol. 60:
17—48.
STEYSKAL, G. C. 1974: Recent advances in the
classification of the Diptera. — Ann.
Entomol. Soc. America 67:513—517.

rufina (Zetterstedt, 1848)
scalaris Meigen, 1830
speciosa Meigen, 1830
triangularis Becker, 1910
troglodytes (Zetterstedt, 1848)
varsoviensis Becker, 1910
Epichlorops Becker, 1910
puncticollis (Zetterstedt, 1848)
Melanum Becker, 1910
laterale (Haliday, 1840)
Diplotoxa Loew, 1863
sg. Diplotoxa Loew, 1863
messoria (Fallén, 1820)
sg. Diplotoxoides Andersson, 1977
dalmatina Strobl, 1900
Cetema Hendel, 1907
cereis (Fallén, 1820)
elongata (Meigen, 1830)
myopina (Loew, 1866)

Thaumatomyia Zenker, 1833
sg. Thaumatomyia Zenker, 1833
hallandica Andersson, 1966
obscurella auctt. nec (Zetterstedt, 1838)
notata (Meigen, 1830)
rufa (Macquart, 1835)
trifasciata (Zetterstedt, 1848)
sg. Chloropisca Loew, 1866
glabra (Meigen, 1830)
Cryptonevra Liroy, 1864
diadema (Meigen, 1830)
tarsata (Fallén, 1820)
Eutropha Loew, 1866
variegata Loew, 1866
Assuania Becker, 1903
distinguenda (Frey, 1909)
Lasiosina Becker, 1910
cinctipes (Meigen, 1830)
parvipennis Duda, 1933
Pseudopachychaeta Strobl, 1902
approximatonervis (Zetterstedt, 1848)

Fam. ACARTOPHTHALMIDAE

Acartophthalmus Czerny, 1902
bicolor Oldenberg, 1910

nigrinus (Zetterstedt, 1848)
pusio Frey, 1947

Fam. CARNIDAE

Carnus Nitzsch, 1818
hemapterus Nitzsch, 1818
Meoneura Rondani, 1856
anceps Frey, 1935
elongella (Zetterstedt, 1838)
flavifacies Collin, 1930
lacteipennis (Fallén, 1823)

lamellata Collin, 1930
neglecta Collin, 1930
neottiophila Collin, 1930
obscurella (Fallén, 1820)
seducta Collin, 1937
triangularis Collin, 1930
vagans (Fallén, 1823)

Fam. MILICHIIDAE

Subfam. Madizinae
Madiza Fallén, 1810
glabra Fallén, 1820
Leptometopa Becker, 1903
latipes (Meigen, 1830)
Neophyllomyza Melander, 1913
acyglossa (Villeneuve, 1920)
Desmometopa Loew, 1865
m-nigrum (Zetterstedt, 1848)
sordidum (Fallén, 1820)

Phylloomyza Fallén, 1810
equitans (Hendel, 1919)
formicae Schmitz, 1923
securicornis Fallén, 1823
silesiaca (Duda, 1935)
tetragona Hendel, 1924
Subfam. Milichiinae
Milichia Meigen, 1830
ludens (Wahlberg, 1847)

Fam. TEPHRITIDAE

Subfam. Ceratitinae
Ceratitis McLeay, 1829
capita (Wiedemann, 1824)
Subfam. Euphrantinae
Euphranta Loew, 1862
connexa (Fabricius, 1794)

Subfam. Trypetinae
Phagocarpus Rondani, 1870
permundus (Harris, 1780)
Cryptaciura Hendel, 1927
rotundiventris (Fallén, 1820)