

*trivialis Loew, Centur. III, 87. — Distr. Columbia.
Prolifica Os., *Lindneri* 4th Report, '70, 1872; also 7th Report.

Observation. About the species of Chlorops enumerated on page 85 of my first Catalogue, Mr. Loew communicates me the following remarks.

- antennalis Fitch, Reports I, 300, see my note [sic] *Agromyza vulgaris*
 annulata Walker, List, etc. IV, 1119. — Huds. B. Terr. [probably Chloropisca. — Loew].
 assimilis Macquart, Dipt. Exot. 4e Suppl. 306, 3; Tab. XXVIII, f. 9. — North America (probably Diplotoxa. — Loew).
 atra Macquart, Dipt. Exot. 4e Suppl. 307, 6; Tab. XXVIII, f. 12 [probably Eutropha; hardly Haplegis. — Loew].
 bistrigata Walker, List, etc. IV, 1120. — Huds. B. Terr. [apparently Chlorops, in the narrower sense. — Loew].
 perlava Walker, List, etc. IV, 1120 [perhaps Diplotoxa. — Loew].
 proxima Say, J. Acad. Phil. VI, 187; Compl. Wr. II, 370. — Indiana.
 soror Macquart, see *Oscinis soror*.
 testacea Macquart, Dipt. Exot. 4e Suppl. 306, 4; Tab. XXVIII, f. 10. — North America [Chlorops, sensu strict. — Loew].
 vittata Fitch, Reports I, 300; Tab. I, f. 5. — New-York. *Oscinis*
 Wiedemann, Auss. Zw. II, 594, 1. — West Indies. [The plumose antennae render the position of this species in the family somewhat doubtful. As *Hippelates eulophus* alone, among all N. A. species, has such antennae, *C. vittata* may be a Hippelates. However South America possesses several Oscinidae with plumose antennae. — Loew.]

Elliponeura.

Loew, Centur VIII, 79; 1869.

*debilis Loew, Centur. VIII, 79. — Distr. Columbia.

Gymnopa.

Fallen, Oscinid., 1820; *Mosillus* Latreille; 1804. (22)

nigroaenea Walker, Dipt. Saund., 413. — United States.
 tarsalis Walker, l. c. — United States.

FAMILY AGROMYZIDAE.

Rhinoessa.

Loew, Wien. Ent. Monatschr. VI, 174. (324)

- *albula Loew, Centur. VIII, 80. — Newport, R. I.
 *coronata Loew, Centur. VI, 98. — Georgia.
 *parvula Loew, Centur. VIII, 81. — Newport, R. I.
Milichia Mels. Syst. Besch. VI, 131; 1830.

Lobioptera.

Wahlberg, Oefvers. af K. Vetensk. Acad. Forh. 1847, 259.

- *arcuata Loew, Zeitschr. f. Ges. Naturw. 1876, 339. — Long Island, N. Y. *Ophthalmomyia*
 *indecora Loew, Centur. VIII, 94. — Nebraska.

placed among
agr. I, 46. —

ve been intro-
eit. f. Entom.;

ad Canada.

ia.

mbia.

ka.

nerica.

1, 318. 1895. Fla.

Phytomyza Fallén, Sp. Ent. Nov. Dipt., 20, 1810.
✓ *securicornis* Fallén, Dipt. Succ. Ochth., 8, 1823.

PHYTOMYZIDAE. — ATEIDAE.

211

Johnson in litt. Oct. 9, 1902.

- ✓ *flaviventris* John. Can. Ent., Sept., 1902,
- ✓ *melampyga* Loew, Centur. VIII, 88. — Distr. Columbia.
- ✓ *neptis* Loew, Centur. VIII, 93. — Nebraska.
- ✓ *parvicornis* Loew, Centur. VIII, 92. — Distr. Columbia.
- ✓ *setosa* Loew, Centur. VIII, 83. — Distr. Columbia.
- ✓ *simplex* Loew, Centur. VIII, 84. — Pennsylvania.
- ✓ *tritici* Fitch, Reports I, 303; Tab. II, f. 1. — New York.
- ✓ *virens* Loew, Centur. VIII, 85. — Pennsylvania.

- picitella* Thomson, Eugen. Resa, 609. — California.
- platyptera* Thomson, Eugen. Resa, 608. — California.

See p. 207.

Cephalomyza Schiner:
Odontocera.

Macquart, Hist. Nat. Dipt. II, 614; 1835. Proc.

- ✓ *dorsalis* Loew, Centur. III, 98. — Distr. Columbia.

Paranyza Will. Kans. Univ. Quart., VI, 1; Jan., 1897.

Phytomyza.

Fallén, Ochthidia; 1823.

- ✓ *nitens* Loew, Centur. VIII, 82. — Pennsylvania.

Oechthiphila.

Fallén, Ochthidia; 1823. (328).

Hispina Thomson, Eugen. Resa, 599. — California.

✓ *polystigma* Meig. Schiner Faun. II, 392, III, var. (Zerk), Wiener Ent. Zeitung, Jan., 1900
Observation. *Ulidia metallica* Bigot, in R. de la Sagra etc. P. 9.

825 belongs to the Agromyzidae, according to Loew, Monogr. III, 202; however in the same volume page 65, he says it may be a

Chrysoomyza, a genus allied to *Ulidia*.

maritima Zett. *Mass.*
juncorum Fall. *Mass.*
elegans Fall. *N.Y.*

FAMILY PHYTOMYZIDAE.

Phytomyza.

Fallén, Sp. Ent. Nov. Dipt., 21, 1810.

Fallén, Phytomyz.; 1823.

- ✓ *clematidis* Loew, Centur. III, 100. — Distr. Columbia.
- ✓ *diminuta* Walker, Trans. Ent. Soc. N. S. IV, 232. — United States.
- ✓ *genualis* Loew, Centur. VIII, 100. — Distr. Columbia. *albiceps* Meig. Becker in litt. to G. H. Hough. *Syn. zetterstedtii* Schin. Same.
- ✓ *ilicicola* Loew, Centur. Vol. II, 290. — Distr. Columbia.
Phytomyza ilicis Loew, Centur. III, 99 (change of name by Loew).
- ✓ *nervosa* Loew, Centur. VIII, 99. — Distr. Columbia.
- solita* Walker, Trans. Ent. Soc. N. Ser. IV, 232. — United States.
- obscura* Fallén, Phytomyz. 4, 8; Meigen, System. Besch. VI, 191;
Staeger, Groenl. Antl. 369, 55. — Europe and Greenland.
- ✓ *chrysanthemii* Kowarz, Linnaeus 7th. Rept., p. 213, 1841.
- ✓ *aguiligis* Hardy. Ann. Ent. Hist., 2nd ser. IV, 392, 1849. Eng. and Fern. Britton in Garden and Forest, Nov. 6, 1895.

FAMILY ASTEIDAE.

Sigaloëssa.

Loew, Centur. VI, 100; 1865. (329).

- ✓ *bicolor* Loew, Centur. VI, 100. — Cuba.

Napomyza Haliday in Westwood's Introd., II, 152; 1840, Type as given, *Phytom. festiva* Meigen.

myz Will.
3, 20.

30.

Förh. 1847,
p. 455.
notations see
North America
ker, Zeit. Hym. Dipt. II, 305, type.

D. Columbia;
— Sweden;
; Fla.; N. Mex.; Col.

Loew in litt.]

ed States.
ab. V, f. 19,

1845. Fla.; Pa.

next Id.