

Musca germinationis, Linn. *Faun. Suec.* 1874. *Gmel. Ed. Syst. Nat.* i. 5, 2857, 122. *Fabr. Ent. Syst.* iv. 356, 183.
Dacus germinationis, *Fabr. Syst. Antl.* 278, 28.

- a. England. Presented by F. Walker, Esq.
- b. England.
- c. France.

OPOMYZA florum, Meig. *Dipt.* vi. 102, 2, *pl.* 59, *f.* 22. *Zett. Ins. Lapp.* 762, 3. *Musca florum*, *Fabr. Ent. Syst.* iv. 352, 164.
Tephritis florum, *Fabr. Syst. Antl.* 317, 3. *Geomyzza punctella*, *Fall. Geom.* 11, 2.

- a. England.

OPOMYZA combinata, var.? Meig. *Dipt.* vi. 109, 18. *Musca combinata?* *Fabr. Ent. Syst.* iv. 352, 163. *Gmel. Ed. Syst. Nat.* i. 5, 2855, 114. *M. germinationis?* *Schrink, Ins. Austr.* 961.
Tephritis combinata? *Fabr. Syst. Antl.* 317, 2. *Geomyzza combinata?* *Fall. Geom.* 2, 1.

- a. France.

OPOMYZA tripunctata, Meig. *Dipt.* vi. 110, 21. *Macq. Hist. Nat. Dipt.* ii. 558, 14. *Geomyzza tripunctata*, *Fall. Geom.* 2, 2. *Zett. Ins. Lapp.* 774, 2.

- a. England.
- b. Upsal. Presented by the Entomological Club.
- c. France.

OPOMYZA maculata, *Macq. Hist. Nat. Dipt.* ii. 558, 15.
a. England. Presented by F. Walker, Esq.

- b. England.

OPOMYZA guttipennis, *Zett. Ins. Lapp.* 762, 4.
a. Lapland. Presented by the Entomological Club.

OPOMYZA apicalis, n. s., fem. *Ferruginea, capite subtus fulvo, abdome supra piceo, palpis antennis pedibusque fulvis, alis limpidis apice fusco maculatis.*

Body pale ferruginous, shining: head and chest beset with very few black bristles: head pale tawny beneath: eyes red: sucker, palpi, and feelers pale tawny: feelers much shorter than the face: third joint conical, longer than the second; bristle slender, black, downy, tawny at the base, more than thrice the length of the third joint: abdomen oval, a little longer than the chest, pitchy above, except at the base: legs tawny; claws black: wings colourless; a brown spot at the tip between the second and third longitudinal

veins; wing-ribs tawny; veins pitchy, tawny towards the base; longitudinal veins straight; lower cross-vein nearly straight, almost straight, parted by more than twice its length from the middle cross-vein, by more than thrice its length from the end of the fourth longitudinal vein, and by less than half its length from the end of the fifth longitudinal vein; poisers whitish. Length of the body $1\frac{1}{2}$ times of the wings 3 lines.

New Zealand. Presented by Dr. Hooker.

OPOMYZA quadrivitta, n. s., fem. *Fulva, thorace vittis quatuor piceis, abdomine vitta fasciisque nigris ornato, palpis antennis pedibusque fulvis, alis subcinereis, costa venisque transversis fusco nebulosis.*

Body tawny: head and chest beset with stout black bristles: head whitish beneath: eyes pitchy: sucker, palpi, and feelers pale tawny: feelers a little shorter than the face: first and second joints set with a few bristles; third conical, longer than the second; bristle black, feathered with hairs on both sides to the tip, about twice the length of the third joint: chest adorned with four pitchy stripes: abdomen oboconical, a little shorter and narrower than the head, thinly clothed with black hairs, beset on each side with a few black bristles, adorned with a black stripe, and on the hind borders of the segments with black bands: legs tawny, beset with a few black hairs and bristles; hind thighs pitchy: wings slightly gray, brown for one-third of the breadth beneath the fore border and at the tip, the outline of the brown is irregular above the cross-veins, which are also clouded with brown; wing-ribs and veins pitchy; longitudinal veins straight; lower cross-vein straight, slightly oblique, parted by much more than its length from the middle cross-vein, by more than thrice its length from the end of the fourth longitudinal vein, and by more than half its length from the end of the fifth longitudinal vein; poisers tawny. Length of the body $1\frac{1}{2}$ line; of the wings 4 lines.

Isle of France.

MILICHIA, Meigen.

MILICHIA speciosa, Meig. *Dipt.* vi. 132, 1, *pl.* 60, *f.* 23—24. Loew, *Ent. Zeit. Stett.* 324.
France.

Walker 1849

